Birdville ISD

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS

School Nurse:	a person who holds a current license as a registered professional nurse (RN) from the Texas Board of Nursing as outlined in the Texas Administrative Code, § 153.1021.
School-Sponsored Activity:	any activity, event, or program occurring on or off school grounds, whether during or outside of regular school hours, which is organized and/or supported by the school
Severe Food Allergy	a dangerous or life-threatening reaction of the human body to a food-borne allergen induced by inhalation, ingestion or skin contact that requires immediate medical attention.

Birdville ISD

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS

Procedure:

1. Notification of a food allergy
   Upon enrollment of a new student, the parent or guardian shall indicate if their child has a known severe food allergy during the online enrollment process.
2. Upon review of the online enrollment records, the campus nurse will:
   identify students with severe food allergies
   as applicable, request parent completes and submits the following forms:

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS

Level III Training: This includes all of Level I and Level II training as well as provides more in-depth training and hands-on training with epi-pen auto injector. The campus Emergency Response Team (ERT) will participate in this level of training as well as other personnel that the school nurse identifies as necessary, such as those employees trained for medication administration.

- More in-depth Level I information
- Identifying students at risk for anaphylaxis
- Planning for students who do not have epinephrine at school
- Development and implementation of FAEAP/IHP/504 plan(s)
- Communication procedures for initiating emergency protocols, including substitute staff.
- Environmental control factors including hand washing and cleaning procedures
- Working with EMS
- Post anaphylaxis debriefing

School Nurse should document Level III training on **Anaphylaxis/Food Allergy Level III/ Epinephrine Documentation** form and **Food Allergy Level 3 Sign In Sheet** and keep in Daily Medication Notebook.

5. In the event of an anaphylactic event, post exposure conference 5 aeposu 13.554 0 Td[f)-28.5(t)-6.6 (he)10.f[an

Severe Food Allergy - Responsibilities of Family & Student

In promoting a safe school environment for children with food allergies at-risk for anaphylaxis, students, parents, and school personnel should work as a team in identifying students at-risk for anaphylaxis, reducing the risk of exposure to food allergens and to react quickly should an exposure or allergic reaction occur.

The responsibilities of the family include:

Notifying the school of the student's allergies in accordance with TEC, Section 25.022. Accurate reporting during the online enrollment process is the preferred method of notification.

Working with the School Nurse (RN) to develop and review the Food Allergy Emergency Action Plan (FAEAP) and Individualized Health Plan (IHP) as well as discuss

Providing properly labeled medications and replace medications after use or upon expiration.

504 accommodations do not apply to before/after school activities provided by an outside agency.

The responsibilities of the student include:

No trading of food with others.

Not eating anything with unknown ingredients or known to contain any allergen.

Being proactive in the care and management of their food allergies and reactions (as developmentally appropriate).

Immediately notifying an adult if they eat something, they believe may contain a food to which they are allergic.

Severe Food Allergy - Responsibilities of Classroom Teacher

In promoting a safe school environment for children with food allergies at-risk for anaphylaxis,

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS

Field Trips:

Give the nurse at least TWO weeks' notice prior to field trips.
Ensure the FAEAP and the student's epinephrine is taken on the field trip and all outings.
Collaborate with parents of student with food allergies when planning field trips.
Consider eating arrangements on field trips and plan for reduction of exposure to a student's life-threatening food allergy.
Consider inviting parents of students at risk for anaphylaxis to accompany their child on school trips, and/or to act as a chaperone. However, the student's safety or attendance must not be a condition on the parent's presence on the trip.
Ensure that 1 or 2 staff members on the field trip are trained in recognizing signs and symptoms of life-threatening allergic reactions and are trained by the school nurse to use an epi-pen.
Consider ways to wash hands and encourage hand washing before and after eating (e.g. provision for hand wipes, etc.).

Severe Food Allergy - Responsibilities of Campus Administrator

In promoting a safe school environment for children with food allergies at-

Birdville ISD

CARE OF THE STUDENT WITH FOOD ALLERGIES AT-RISK FOR ANAPHYLAXIS